

# Interaction of cyanine dyes with nucleic acids.

## 6. Synthesis and spectroscopic properties of thiazole orange—amino acids conjugates

Sergiy M. Yarmoluk, Dmytro V. Kryvorotenko, Yuriy S. Gerasymchuk

Institute of Molecular Biology and Genetics, National Academy of Sciences of Ukraine  
150 Acad. Zabolotnoho vul., Kyiv, 252143, Ukraine

*The synthesis of amino acid (L-Trp and L-Tyr) derivatives of thiazole orange, monomethine cyanine dye which significantly increases fluorescence intensity when bound to nucleic acids is described. Hydroxysuccinimide ester of Cyan 6 (2-[4-N-Carboxyethyl-quinoline)-methyl]-3-methylbenzothiazole-1,4-p-toluenesulfonate) was used for the conjugation of dye with amino acids. Interaction of obtained conjugates with nucleic acids was investigated using spectroscopic methods.*

**Materials and Methods.** The TLC was performed on Kieselgel 60 F<sub>254</sub> plates («Merck», Germany) using CHCl<sub>3</sub>—MeOH—AcOH (8.5:1.0:0.5) system. FAB MS analyses were carried out in the positive mode using mass spectrometer MI-1201E (PO «Elektron», Ukraine).

**Spectroscopic measurements.** The absorption spectra were recorded on «Specord M-40» (VEB Carl Zeiss Jena, Germany). Fluorescence spectra were obtained with fluorescence spectrophotometer Hitachi Model 850 (Japan). Fluorescence was excited with 150W Xe-lamp emission and measurements were carried out in thermostatable quartz cell (0.5 × 0.5 cm). All spectra were corrected by multiplying fluorescence intensities measured over an interval of 5 nm by proper correction factor for corresponding wavelengths. In corrected spectra fluorescence intensity values were proportional to a numbers of photons per unit of wavelength interval.

**Preparation of DNA, RNA and dyes stock solutions.** Stock solutions of dyes (2 · 10<sup>-3</sup> M) were prepared by dissolving dyes in DMSO. All dyes were stable under these conditions for several months, whereas in aqueous solutions some dyes gradually lost their fluorescence properties. Working solutions were prepared immediately prior to use. For spectral studies total calf thymus DNA («Sigma») and yeast RNA

(«Sigma») were used. Nucleic acids stock solutions were prepared in TE buffer (Tris-HCl, 50 mM, EDTA, 10 mM, pH 8.0) in concentration 6 · 10<sup>-3</sup> b. p./ml for DNA and 1.2 · 10<sup>-2</sup> b. p./ml for RNA.

**Absorbance and fluorescent emission spectra.** For spectral measurements the complexes of dyes with nucleic acids were obtained in Tris-HCl (50 mM, pH 8.0) buffer by mixing of dye stock solution with DNA or RNA solution. The final concentration of NA was 10<sup>-4</sup> and 2.0 · 10<sup>-4</sup> mM respectively. Final dyes concentrations were 0.02 mM. Dye—nucleic acid complexes were prepared with approxymal ratio 1 dye per 20 b. p. of RNA and per 10 b. p. of DNA. For optical measurement of free cyanines the same dyes concentrations were used.

**Fluorescence titration of Cyan 6—Trp and Cyan 6—Tyr** were carried out in 0.05 M Tris buffer (pH 8) with total calf thymus DNA and yeast RNA at 540 nm for DNA and 545 nm for RNA. Fixed concentration of DNA 6 · 10<sup>-3</sup> M and 1.2 · 10<sup>-2</sup> M were used for the titration of fixed concentration of dye (Fig. 3). The dye concentration was 10<sup>-5</sup> M. The DNA and RNA concentration was changed from 2.5 · 10<sup>-6</sup> to 10<sup>-4</sup> from 5 · 10<sup>-6</sup> to 2 · 10<sup>-4</sup> respectively.

**General procedure for the synthesis cyanine dye—amino acids conjugates.** The succinimidyl ester of Cyan 6 was prepared by dissolving the dye (5.34 mg, 0.01 mmol) in 1 ml of dry DMFA and then adding N-hydroxysuccinimide (1.15 mg, 0.01 mmol) and




Fig. 1. Synthetic scheme of preparation of dye—amino acid conjugates

DCC (4.12 mg, 0.02 mmol) to stirred solution at 0 °C. Mixture was stirred for 4 h. Then amino acid ester hydrochloride (0.011 mM) and 0.05 ml of dry pyridine were added. Reaction mixture was stirred for 2 h at 0 °C and was kept in refrigerator for 18 h. Purification of amino acid—Cyan 6 conjugates was performed on the prepared C3 stationary phase [1]. Conjugates were recovered from the fractions with a rotary evaporator at 40 °C.

*Cyan 6—Trp—OCH<sub>3</sub>*.  $R_f = 0.67$  MS:m/e (relative intensity) 563 (MH<sup>+</sup>, 5), 363 (14), 317 (3), 290 (6), 275 (5), 264 (4), 219 (4), 150 (16).

*Cyan 6—Tyr—OCH<sub>3</sub>*.  $R_f = 0.72$  MS:m/e (relative intensity) 540 (MH<sup>+</sup>, 14), 526 (6), 407(7), 363 (27), 317 (7), 303 (7), 290 (12), 275 (9), 219 (6), 150 (18).

**Results and Discussion.** The cyanine monomethine dyes have many of properties presented as ideal for the fluorescent homogeneous detection of nucleic acids. With this determination principle no separation of unbound probe is necessary [2]. A series of new monomeric and dimeric fluorescent cyanines have recently been developed for this purpose [3, 4]. Monomethine cyanine dyes seem to possess all characteristics for quantitation of unknown nucleic acids in solutions, as low intrinsic fluorescence of unbound probe, large fluorescence enhancement upon probe binding, large linear detection range, and sequence non-specific detection [5].

The main research of our laboratory concerns development of sequence specific homogeneous detection of nucleic acids in solution. Recently pyrylium cyanine dye — oligonucleotide conjugate was used for

described by Waggoner et al. [7]. Functional groups in proposed cyanine reagents were sulfhydryl [7], isothiocyanate [8] and succinimidyl esters [9]. Procedure using active N-hydroxysuccinimide esters of dye seems to be more convenient for their conjugation with  $\alpha$ -amino group of amino acids or peptides. Besides we have shown recently that incorporation of carboxylic acid groups into basic cyanine structure of TO did not influence on fluorescence probe properties [10] (Fig. 1).

Succinimidyl ester of dye carboxyl group and methyl esters of hydrochlorides of amino acids in pyridine were used to synthesize amino acid—dye conjugates. Active esters were prepared with use of DCC and hydroxysuccinimide (HONSu) in DMFA.

Recently published methods for the HPLC of cyanine dyes use C<sub>18</sub> stationary phase [11]. Unfortunately, used cyanine dyes are very hydrophobic and positively charged compounds. As a result, they have large retention time on HPLC stationary phases. In order to overcome these problems TMS-silica (C<sub>3</sub>) stationary phase prepared according to [1] was successfully used (Fig. 2). Chemical structures of synthesized conjugates were confirmed by fast atom bombardment mass spectrometric (FAB MS) analyses.

Spectroscopic data for two synthesized cyanine conjugates and Cyan 6 (TO analog) are presented in Table. 1. Dye Cyan 6 with previously investigated spectral properties was included for the comparison [9]. Absorption spectra of Cyan 6 and its amino acid




Table 1  
Chemical structures and spectroscopic properties of cyanine dyes

Dye	Structure	$abs_{\lambda_{max}}$ DMFA [buff.], nm	$em_{\lambda_{max}}$ , nm	$\Delta S$ , nm	$F^*$
Cyan 6-Trp	<p>CH<sub>3</sub>      CF<sub>3</sub> COO-</p>	507 [475, 505]	**	-	0.076
Cyan 6-Tyr	<p>CH<sub>3</sub>      CF<sub>3</sub>COO-</p>	506 [470, 506]	-	-	0.062
Cyan 6	<p>CH<sub>3</sub>      OTos<sup>-</sup></p>	503 [475, 506]	-	-	0.03

\*Arbitrary units; \*\*There is no clear maxima.

derivatives were similar. The wavelengths of absorption maxima ( $abs_{\lambda_{max}}$ ) of Cyan 6-Trp and Cyan 6-Tyr slightly depended on solvent. Cyanine dyes showed the red shift (5--7 nm) after going from aqueous buffer to less polar DMSO.

The fluorescence of conjugates and free Cyan 6 were very low and fluorescence spectra had no clear maxima. Fluorescence intensity of amino acids derivatives were slightly higher as compared to free dye. The data on the absorbance and fluorescence

Table 2  
Spectroscopic properties of cyanine dye complexes with nucleic acids

Dye	DNA—Dye complex					RNA—Dye complex				
	abs. $\lambda_{max}$ , nm	em. $\lambda_{max}$ , nm	$\Delta S$ , nm	$I$	$\Delta Q$	abs. $\lambda_{max}$ , nm	em. $\lambda_{max}$ , nm	$\Delta S$ , nm	$I$	$\Delta Q$
Cyan 6—Trp	478.5	540	32	28.8	380	478.5	545	40	22.2	292
Cyan 6—Trp	479.5	540	32	24.5	395	478.5	543	38	21.1	324
Cyan 6	478.5	540	32	7.1	237	478.5	545	39	13.5	450




Fig. 3. Fluorescence titration of Cyan 6—Trp (1) and Cyan 6—Tyr (2) with DNA (A) and RNA (B). Fluorescence intensity is in arbitrary units.

emission spectra of the nucleic acid—bound form of conjugates are presented in Table 2. The fluorescent intensity of DNA complexes of amino acids derivatives increased 1.6 times comparing with similar complex of Cyan 6. Fluorescent intensity of RNA complexes decreased 1.5 times for Trp derivative and 1.3 times for Tyr derivative in contrast to free Cyan 6 complex. All dyes had medium Stokes shifts (32 nm for DNA complexes and 38—40 nm for RNA complexes).

Data on fluorescence titration of Cyan 6—Trp and Cyan 6—Tyr are presented in Fig. 3. For the Cyan 6—Trp—DNA complexes fluorescence plateau is reached at 4 base pairs per dye (A). Titration of RNA showed that the fluorescence emission of bound Cyan 6—Trp reaches a maximum at 15 base/dye (B).

New amino acid derivatives of cyanine dyes were first synthesized with use of active N-hydroxysuccinimide esters of dyes.

Acknowledgements. The authors gratefully ac-

knowledge the assistance of Ms. Vladislava Kovalska in spectral characterisation of dye—amino acid conjugates. We thank Dr. Yury Kovtun for providing us with the sample of cyanine dye Cyan 6.

*С. М. Ярмолюк, Д. В. Криворотенко, Ю. С. Герасимчук*

Взаємодія ціанінових барвників з нуклеїновими кислотами. 6. Синтез та спектроскопічні властивості кон'югатів тіазолового оранжевого з амінокислотами

Резюме

Описано синтез амінокислотних похідних (*L*-Trp та *L*-Tyr) з тіазоловим оранжевим — монометиновим ціаніновим барвником, що значно підвищує інтенсивність власної флюоресценції при зв'язуванні з нуклеїновими кислотами. Гідроксисукцинімідний ефір Cyan 6 (2-[*N*-карбокситіл-хінолін)-метил]-3-метилбензтіазол-1,4-*p*-толуолсульфата) був використаний для кон'югації. Взаємодію синтезованих кон'югатів з нуклеїновими кислотами досліджено спектроскопічними методами.

*С. Н. Ярмолюк, Д. В. Криворотенко, Ю. С. Герасимчук*

Взаимодействие цианиновых красителей с нуклеиновыми кислотами. 6. Синтез и спектроскопические свойства конъюгатов тиазолового оранжевого с аминокислотами

Резюме

Описан синтез аминокислотных производных (*L*-Trp и *L*-Tyr) с тиазоловым оранжевым — монометиновым цианиновым красителем, который значительно увеличивает интенсивность собственной флюоресценции при связывании с нуклеиновыми кислотами. Гидроксисукцинимидный эфир Cyan 6 (2-[*N*-карбокситил-хинолин)-метил]-3-метилбензтиазол-1,4-*p*-толуолсульфата) был использован для конъюгации красителя с аминокислотами. Взаимодействие синтезированных конъюгатов с нуклеиновыми кислотами исследовано спектроскопическими методами.

#### REFERENCES

1. Калашиников В. В., Самуков В. А., Шубина Т. Н. Использование обращенно-фазовой хроматографии в олигонуклеотидном синтезе // Биоорг. химия.—1983.—9.—С. 666—672.

2. Schwartz H. E., Ulfelder K. J. Capillary electrophoresis with laser-induced fluorescence detection of PCR fragments using thiazole orange // Anal. Chem.—1992.—64, N 15.—P. 1737—1740.
3. Haughland R. P. Handbook of fluorescent probes and research chemicals // Molecular Probes.—Eugene: OR, 1996.—679 p.
4. Yarmoluk S. M., Kovalska V. B., Smirnova T. V., Shandura M. P., Kovtun Yu. P., Matsuka G. Kh. Interaction of cyanine dyes with nucleic acids. 2. Spectroscopic properties of methyleneoxy analogues of thiazole orange // Biopolimery i kletka.—1996.—12, N 6.—P. P. 74—81.
5. Skeidsvoll J., Ueland P. M. Analysis of double-stranded DNA by capillary electrophoresis with laser-induced fluorescence detection using the monomeric dye SYBR Green I // Anal. Biochem.—1995.—231—P. 359—365.
6. Yarmoluk S. M., Kostenko A. M., Kornishyna O. S., Dubey I. Y. Interaction of cyanine dyes with nucleic acids. 4. Efficient 5'-fluorescent labelling of oligonucleotides with monomethylene pyrylium cyanine dye, Cyan 39 // Biopolimery i kletka.—1998.—14, N 1.—P. 82—86.
7. Ernst L. A., Gupta R. K., Mujumdar R. B., Waggoner A. S. Cyanine dye labelling reagents for sulfhydryl groups // Cytometry.—1989.—10.—P. 3—10.
8. Mujumdar R. B., Ernst L. A., Mujumdar S. R., Waggoner A. S. Cyanine dye labelling reagents containing isothiocyanate groups // Cytometry.—1989.—10.—P. 11—19.
9. Southwick P. L., Ernst L. A., Tauriello E. W., Stephen R. P., Mujumdar R. B., Mujumdar S. R., Clever U. A., Waggoner A. S. Cyanine dye labelling reagents — carboxymethylindocyanine succinimidyl esters // Cytometry.—1990.—11.—P. 418—430.
10. Yarmoluk S. M., Zhyvoloup A. N., Koval'ska, Klimenko I. V., Slomisky Yu. L. Interaction of cyanine dyes with nucleic acids. I. Studies on monomethylene cyanine dyes as possible fluorescent probes for the detection of nucleic acids // Biopolimery i kletka.—1996.—12.—P. 69—74.
11. Akins D. L., Kurnar V. T. High performance liquid chromatography of cyanine dyes Multiphase separation, purification, and substitution of the counter ion // J. Chrom. A.—1995.—689.—P. 269—273.

УДК 342.95  
Received 25.02.98